

Mean field ground state of a spin-1 condensate in a magnetic field

Wenxian Zhang, Su Yi and Li You

School of Physics, Georgia Institute of Technology, Atlanta GA 30332-0430, USA

Abstract. We revisit the topic of the mean field ground state of a spin-1 atomic condensate inside a uniform magnetic field (B) under the constraints that both the total number of atoms (N) and the magnetization ($\mathcal{M}$) are conserved. In the presence of an internal state (spin component) independent trap, we also investigate the dependence of the so-called single spatial mode approximation (SMA) on the magnitude of the magnetic field and $\mathcal{M}$. Our result indicate that the quadratic Zeeman effect is an important factor in balancing the mean field energy from elastic atom-atom collisions that are known to conserve both N and $\mathcal{M}$.

1. Introduction

Atomic Bose-Einstein condensates (BEC) have provided a successful testing ground for theoretical studies of quantum many-body systems [1]. In earlier BEC experiments, atoms were spatially confined with magnetic traps, which essentially freeze the atomic internal degrees of freedom [2]. Most studies were thus focused on scalar models, i.e. single component quantum degenerate gases [3]. More recently, the emergence of spin-1 condensates [4, 5, 6] (of atoms with hyperfine quantum number $F = 1$) has created opportunities for understanding degenerate gases with internal degrees of freedom [7, 8, 9, 10, 11, 14].

In this paper, we investigate the mean field ground state structures of a spin-1 atomic condensate in the presence of an external magnetic field (B). We focus on several aspects of the ground state properties strongly affected by the requirement that elastic atom-atom collisions conserve both the total number of atoms (N) and the magnetization ($\mathcal{M}$). Several earlier studies have focused on the global ground state structures when the conservation of $\mathcal{M}$ was ignored, or in the limiting case of a vanishingly small magnetic field ($B = 0$) [6, 7, 8, 9, 10, 11, 14]. As we show in this study, in the presence of a nonzero magnetic field, the conservation of $\mathcal{M}$ leads to ground state population distributions significantly different from that of the global ground state.

Our system is described by the Hamiltonian (repeated indices are summed) [7]

$$\mathcal{H} = \int d\vec{r} \psi_i^\dagger (\mathcal{L}_{ij} + H_{ZM}) \psi_j + \frac{c_0}{2} \int d\vec{r} \psi_i^\dagger \psi_j^\dagger \psi_j \psi_i + \frac{c_2}{2} \int d\vec{r} \psi_k^\dagger \psi_i^\dagger (F_\eta)_{ij} (F_\eta)_{kl} \psi_j \psi_l, \quad (1)$$

where $\psi_j(\vec{r})$ is the field operator that annihilates an atom in the j -th ($j = +, 0, -$) internal state at location $\vec{r}$, $\mathcal{L}_{ij} \equiv [-\hbar^2 \nabla^2 / 2M + V_{\text{ext}}(\vec{r})] \delta_{ij}$ with M the mass of each atom and $V_{\text{ext}}(\vec{r})$ an internal state independent trap potential. Terms with coefficients c_0 and c_2 of Eq. (1) describe elastic collisions of the spin-1 atom ($|F = 1, M_F = +, 0, -\rangle$), expressed in terms of the scattering length a_0 (a_2) for two spin-1 atoms in the combined symmetric channel of total spin 0 (2), $c_0 = 4\pi\hbar^2(a_0 + 2a_2)/3M$ and $c_2 = 4\pi\hbar^2(a_2 - a_0)/3M$. $F_{\eta=x,y,z}$ are spin-1 matrices with

$$F_x = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad F_y = \frac{i}{\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad F_z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

The external magnetic field B is taken to be along the quantization axis ($\hat{z}$), it induces a Zeeman shift on each atom given by

$$H_{ZM}(B) = \begin{pmatrix} E_+ & 0 & 0 \\ 0 & E_0 & 0 \\ 0 & 0 & E_- \end{pmatrix}.$$

According to the Breit-Rabi formula [12], the individual level shift can be expressed as

$$E_+ = -\frac{E_{\text{HFS}}}{8} - g_I \mu_I B - \frac{1}{2} E_{\text{HFS}} \sqrt{1 + \alpha + \alpha^2},$$

$$\begin{aligned}
E_0 &= -\frac{E_{\text{HFS}}}{8} - \frac{1}{2}E_{\text{HFS}}\sqrt{1+\alpha^2}, \\
E_- &= -\frac{E_{\text{HFS}}}{8} + g_I\mu_I B - \frac{1}{2}E_{\text{HFS}}\sqrt{1-\alpha+\alpha^2},
\end{aligned} \tag{2}$$

where E_{HFS} is the hyperfine splitting [12], and g_I is the Lande g -factor for the atomic nuclei with nuclear spin $\vec{I}$. μ_I is the nuclear magneton and $\alpha = (g_I\mu_I B + g_J\mu_B B)/E_{\text{HFS}}$ with g_J the Lande g -factor for the valence electron with total angular momentum $\vec{J}$. μ_B is the Bohr magneton.

2. Mean field approximation

At near zero temperatures and when the total number of condensed atoms is large, the ground state is essentially determined by the mean field term $\Phi_i = \langle\psi_i\rangle$. Neglecting all quantum fluctuations we arrive at the mean field energy functional from Eq. (1) [8, 13]

$$H[\{\Phi_i\}] = H_S + E_0 N + \frac{c_2}{2}\langle\vec{F}\rangle^2 - \eta_0\langle F_z\rangle + \delta\langle F_z^2\rangle, \tag{3}$$

where the symmetric part

$$H_S = \int d\vec{r} \left[\Phi_i^* \mathcal{L}_{ij} \Phi_j + \frac{c_0}{2} \Phi_i^* \Phi_j^* \Phi_j \Phi_i \right], \tag{4}$$

is invariant under the exchange of spin component indices, thus is independent of the external B field. The Zeeman shift as given by the Breit-Rabi formula (2) can be described by two positive parameters [13]

$$\begin{aligned}
2\eta_0 &= E_- - E_+, \\
2\delta &= E_+ + E_- - 2E_0,
\end{aligned} \tag{5}$$

which measure approximately the linear and quadratic Zeeman effects. The B -field dependence of η_0 and δ for a ^{87}Rb atom are displayed in Fig. 1.

The elastic atomic collisions as described by the c_0 and c_2 parts of the Hamiltonian (1) conserve both N and $\mathcal{M}$, which in the mean field approximation are given by

$$\begin{aligned}
N &= \sum_{j=\pm,0} \int d\vec{r} \langle \psi_j^\dagger(\vec{r}) \psi_j(\vec{r}) \rangle \approx \sum_{j=\pm,0} \int d\vec{r} |\Phi_j(\vec{r})|^2, \\
\mathcal{M} &= \int d\vec{r} [\langle \psi_+^\dagger(\vec{r}) \psi_+(\vec{r}) \rangle - \langle \psi_-^\dagger(\vec{r}) \psi_-(\vec{r}) \rangle] \\
&\approx \int d\vec{r} [|\Phi_+(\vec{r})|^2 - |\Phi_-(\vec{r})|^2].
\end{aligned} \tag{6}$$

Before continuing our discussion of the ground state structures, we shall first briefly comment on the importance of the above two constraints. In a typical experiment, the last stage before condensation consists of atomic evaporations, during which neither N nor $\mathcal{M}$ is conserved. For a scalar condensate, typically the ground state is obtained from a minimization of Eq. (3) subjected to the constraint of only N conservation. This gives rise to the Gross-Pitaevskii equation (GPE) and the associated condensate chemical potential, which mathematically is simply the Lagrange multiplier of the constrained




Figure 1. Approximate linear and quadratic Zeeman effects as characterized by parameters η_0 and δ versus magnetic field B for a ^{87}Rb atom.

minimization. A spin-1 condensate requires the introduction of two Lagrange multipliers during the minimization subjected to both the N and $\mathcal{M}$ conservation constraints, as was first performed in [13].

When atomic interactions are ferromagnetic ($c_2 < 0$ as for ^{87}Rb atoms) and when the external B -field is negligible, we have shown previously that the ground state structure is simply a state where all individual atomic spins are aligned in the same direction [14]. In this case, the conservation of $\mathcal{M}$ can be simply satisfied by tilting the quantization axis away from the direction of the condensate spin. This can always be done if a system described by (1) is rotationally symmetric, and thus contains the $\text{SO}(3)$ symmetry [7]. The presence of a nonzero B field, on the other hand, breaks the rotational symmetry, [e.g. the linear Zeeman shift, reduces the $\text{SO}(3)$ to $\text{SO}(2)$ symmetry], thus the conservation of $\mathcal{M}$ has to be included in the minimization process directly.

The global ground state phase diagram including both linear and quadratic Zeeman effect was first investigated by Stenger *et al* [6]. In this early study, although the $\mathcal{M}$ conservation was included in their formulation, it was not separately discussed, consequently their results do not easily apply to systems with fixed values of $\mathcal{M}$. The ground state structures as given in Ref. [6] correspond to the actual ground state as realized through a $\mathcal{M}$ non-conserving evaporation process (e.g. in the presence of a nonzero B -field) that serves as a reservoir for condensate magnetization. Our study to be presented here, on the other hand, would explicitly discuss the phase diagram for fixed values of $\mathcal{M}$, which could physically correspond to experimental ground states (with/without a B -field) due to a $\mathcal{M}$ conserving evaporation process. Although more limited, as our results can be traced to linear trajectories of $\mathcal{M} = \text{const.}$ in the phase diagram of Ref. [13], we expect them to be useful, especially in predicting ground state structures when a ready-made spinor condensate is subjected to external manipulations

that conserve both N and $\mathcal{M}$.

When atomic interactions are anti-ferromagnetic ($c_2 > 0$), the global ground state was first determined to be a total spin singlet [8]. More elaborate studies, including quantum fluctuations, were performed by Ho and Yip [17] and Koashi and Ueda [18]. Unfortunately, these results [17, 18] do not correspond to actual ground states as realized in current experiments, because of the presence of background magnetic fields. For instance, the states as found in Ref. [17] are only possible if the magnetic field B is less than $70\mu\text{G}$ at the condensate density as realized in the MIT experiments [13]. The Zeeman shift (see Fig. 1) due to the presence of even a small magnetic field can overwhelm atomic mean field interaction and typical atomic thermal energy, thus if it were not for the conservation of $\mathcal{M}$, the ground state would simply correspond to all atoms condense into the lowest Zeeman sublevel of $|M_F = 1\rangle$.

We now minimize H of Eq. (3) by denoting $\Phi_j(\vec{r}) = \sqrt{N_j}\phi_j(\vec{r})e^{-i\theta_j}$, with a real mode function $\phi_j(\vec{r})$ ($\int \phi_j^2(\vec{r})d\vec{r} = 1$) and phase θ_j . It is easy to check that the phase convention of ferromagnetic/anti-ferromagnetic interactions as obtained previously [10] in the absence of a B -field still remains true, i.e.

$$\theta_+ + \theta_- - 2\theta_0 = 0, \quad c_2 < 0 \text{ (ferromagnetic)}, \quad (7)$$

$$\theta_+ + \theta_- - 2\theta_0 = \pi, \quad c_2 > 0 \text{ (anti-ferromagnetic)}. \quad (8)$$

3. Ground state in a homogeneous system

In a homogeneous system such as a box type trap (of volume $\mathcal{V}$), adopting the above phase convention, the resulting ground state energy functional becomes (+/− for $c_2 < 0$ and $c_2 > 0$ respectively)

$$H[\{N_i\}] = H_S + E_0N + \frac{c_2}{2\mathcal{V}} \left[(N_+ - N_-)^2 + 2N_0(\sqrt{N_+} \pm \sqrt{N_-})^2 \right] - \eta_0(N_+ - N_-) + \delta(N_+ + N_-). \quad (9)$$

Expressing everything in terms of fractional populations and fractional magnetization $n_i = N_i/N$ and $m = \mathcal{M}/N$, and note that $n_+ + n_- = 1 - n_0$, $n_+ - n_- = m$, Equation (9) becomes

$$\frac{H[\{n_i\}]}{N} = \frac{H_S}{N} + E_0 + c \left[(n_+ - n_-)^2 + 2n_0(\sqrt{n_+} \pm \sqrt{n_-})^2 \right] - \eta_0(n_+ - n_-) + \delta(n_+ + n_-), \quad (10)$$

with an interaction coefficient $c = c_2N/2\mathcal{V}$, tunable through a change of condensate density.

We now minimize Eq. (10) under the two constraints $n_+ + n_0 + n_- = 1$ and $n_+ - n_- = m$. We restrict our discussion to the region $-1 < m < 1$ as the special cases of $m = \pm 1$ are trivial. Because H_S , E_0 , c , η_0 , and m are all constants for given values of B , N , and $\mathcal{V}$, the only part left to be minimized is

$$\mathcal{F} = 2cn_0(\sqrt{n_+} \pm \sqrt{n_-})^2 + \delta(n_+ + n_-). \quad (11)$$

In the special case of $c = 0$, Equation (11) reduces to

$$\mathcal{F} = \delta(n_+ + n_-). \quad (12)$$

The ground state is then very simple. When $\delta > 0$, which seems to be always the case for quadratic Zeeman shift, the minimum is reached by having as large a n_0 (thus as small a $n_+ + n_-$) as possible, namely

$$n_0 = 1 - |m|, \quad n_+ = \begin{cases} |m|, & m \geq 0 \\ 0, & m < 0 \end{cases}, \quad n_- = \begin{cases} 0, & m \geq 0 \\ |m|, & m < 0 \end{cases} \quad (13)$$

When $\delta = 0$, we have (in general) three condensate components with $n_{\pm} = (1 - n_0 \pm m)/2$ and $0 \leq n_0 \leq 1 - |m|$.

For ferromagnetic interactions with $c < 0$, we define $x = n_+ + n_-$. The ground state is then determined by the minimum of

$$\mathcal{F} = g_+(x) + \delta x, \quad (14)$$

with $g_+(x) \equiv 2c(1 - x)(x + \sqrt{x^2 - m^2})$. When $\delta = 0$, we find

$$n_{\pm} = \frac{1}{4}(1 \pm m)^2, \quad n_0 = \frac{1}{2}(1 - m^2), \quad (15)$$

which is the same as obtained in [11, 14]. However with a nonzero $\delta > 0$, we find in general

$$n_{\pm} = \frac{1}{2}(x_0 \pm m), \quad n_0 = 1 - x_0 \geq \frac{1}{2}(1 - m^2), \quad (16)$$

with x_0 being the root of equation $g'_+(x) + \delta = 0$, it turns out that there always exists one and only one solution to the equation. The equilibrium value for n_0 is larger than the result of Eq. (15) because the quadratic Zeeman effect causes a lowering of the total energy if two $|M_F = 0\rangle$ atoms are created when an $|M_F = +1\rangle$ atom collides with an $|M_F = -1\rangle$ atom. Figure 2 displays the results of Eq. (16) for a typical ^{87}Rb condensate, for which the atomic parameters are $E_{\text{HFS}} = (2\pi)6.8347\text{GHz}$ [12], $a_0 = 101.8a_B$, and $a_2 = 100.4a_B$ (a_B is the Bohr radius) [15]. At weak magnetic fields, typically a condensate contains all three spin components. With the increasing of B -field, the quadratic Zeeman effect becomes important which energetically favors the $|0\rangle$ component, so typically only two components survive: the $|0\rangle$ component and the larger (initial population) of the $|+\rangle$ or $|-\rangle$ component, so the ground state becomes (for $m > 0$) $n_+ \simeq m$ and $n_0 \simeq 1 - m$.

Finally we consider the case of anti-ferromagnetic interactions for $c > 0$, we have then

$$\mathcal{F} = g_-(x) + \delta x, \quad (17)$$

with $g_-(x) = 2c(1 - x)(x - \sqrt{x^2 - m^2})$. For $\delta = 0$, we again recover the standard result

$$n_0 = 0, \quad n_{\pm} = \frac{1}{2}(1 \pm m), \quad (18)$$

if $m \neq 0$. When $m = 0$, the ground state is under-determined as many solutions are allowed as long as they satisfy $n_+ = n_- = 1 - n_0$ with $n_0 \in [0, 1]$.




Figure 2. The dependence of fractional population for different spin component on m and B for a spin-1 ^{87}Rb homogeneous condensate with $N/V = 5 \times 10^{14} \text{cm}^{-3}$.




Figure 3. The same as in Fig. 2, but now for a spin-1 ^{23}Na condensate.

In an external B -field when $\delta > 0$, we first consider the special case of $m = 0$. It can be easily seen from Eq. (17) that $n_0 = 1$ is the ground state. For $m \neq 0$, we obtain the following result: when $\delta > 2c[1 - \sqrt{1 - m^2}]$, the ground state will have three condensate components with

$$n_{\pm} = (x_0 \pm m)/2, \quad n_0 = 1 - x_0, \quad (19)$$

where x_0 is the root of equation $g'_-(x) + \delta = 0$; When $\delta \leq 2c[1 - \sqrt{1 - m^2}]$, only $|+\rangle$ and $|-\rangle$ components exist, i.e., $n_{\pm} = (1 \pm m)/2$.

Figure 3 is the typical results for a spin-1 ^{23}Na condensate. The atomic parameters are $E_{\text{HFS}} = (2\pi)1.7716\text{GHz}$ [12], $a_0 = 50a_B$ and $a_2 = 55a_B$ [16]. At $B = 0$ there are only two condensate components, $|+\rangle$ and $|-\rangle$. For $B > 0$ but not very strong, there are two possibilities: three nonzero condensate components if $m < m_c$ and two nonzero condensate components if $m \geq m_c$, with $\delta(B) = 2c(1 - \sqrt{1 - m_c^2})$. When B -field gets stronger, i.e. $\delta(B) \geq 2c$, there are always three condensate components.

Figure 4 summarizes the ground state structures of a homogeneous spin-1 condensate in a B -field for different c and m .




Figure 4. The ground state phase diagram for a homogeneous spin-1 condensate. Dashed curves and lines denote gradual transitions across the boundaries, solid lines denote discontinuous jumps. x_0 is the solution to equation $g'_\pm(x) + \delta = 0$ and the curves for $c > 0$ is determined by $\delta(B) = 2c[1 - \sqrt{1 - m^2}]$. The open circle at $B = 0, m = 0$ for $c > 0$ denotes the family of degenerate ground state $(\frac{1-n_0}{2}, n_0, \frac{1-n_0}{2})$.

4. Ground state inside a harmonic trap

In the previous section, we investigated in detail mean field ground state structures for a spin-1 condensate in a homogeneous confinement. For the case of a harmonic trap as in most experiments, there is no reason to believe *a priori* that the above conclusions still hold. In fact, the structures and phase diagrams as discussed before is only meaningful if the spatial mode function $\phi_j(\vec{r})$ for different spin components is identical. Otherwise, it would be impossible to classify the rich variety of possible solutions. When the spatial mode functions are the same, the spatial confinement simply introduces an average over the inhomogeneous density profile of the mode function.

The aim of this section, is therefore to determine the validity of the single mode approximation (SMA) in the presence of an external B -field and a harmonic trap. For simplicity, we assume the trap to be spherically symmetric. We employ numerical methods to directly find the ground state solutions from the coupled Gross-Pitaevskii equation

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} \Phi_+ &= [\mathcal{H} + E_+ - \eta + c_2(n_+ + n_0 - n_-)]\Phi_+ + c_2\Phi_0^2\Phi_-^*, \\
 i\hbar \frac{\partial}{\partial t} \Phi_0 &= [\mathcal{H} + E_0 + c_2(n_+ + n_-)]\Phi_0 + 2c_2\Phi_0^*\Phi_+\Phi_-, \\
 i\hbar \frac{\partial}{\partial t} \Phi_- &= [\mathcal{H} + E_- + \eta + c_2(n_- + n_0 - n_+)]\Phi_- + c_2\Phi_0^2\Phi_+^*,
 \end{aligned} \tag{20}$$

subjected to the conservations of both N and $\mathcal{M}$ [Eqs. (6)]. $\mathcal{H} = -\hbar^2\nabla^2/2M + V_t(\vec{r}) + c_0n$, $n_j = |\Phi_j|^2$, $V_t(\vec{r}) = M\omega^2r^2/2$, and $n = n_+ + n_0 + n_-$. η is the Lagrange multiplier introduced to numerically enable the conservation of $\mathcal{M}$.

It was shown previously that in the absence of an external B -field, and for ferromagnetic interactions, the SMA is rigorously valid despite the presence of a harmonic trap [14]. We can also show that in the presence of a nonzero B -field, the linear Zeeman shift does not affect the validity of the SMA because it can be simply




Figure 5. Typical densities of spatial mode functions for each components of a ^{87}Rb (a) and a ^{23}Na (b) condensate. The solid line denotes the $|+\rangle$ component, the dashed line the $|-\rangle$ component, and the dash-dotted line the $|0\rangle$ component. The parameters are, $N = 10^6$, $\omega = 2\pi \times 100$ (Hz), $B = 1.0$ (Gauss), and $m = 0.5$.

balanced by the external Lagrange multiplier η . The quadratic Zeeman effect, on the other hand, can not be simply balanced, as it favors the production of two $|0\rangle$ atoms by annihilating one $|+\rangle$ and one $|-\rangle$ atom during a collision. Such unbalanced elastic collisions thus break the $\text{SO}(3)$ symmetry of the freedom for an arbitrary quantization axis. Therefore, we do not in general expect the SMA to remain valid inside a nonzero B -field.

Numerically, we find the ground state solutions of Eq. (20) by propagating the equations in imaginary time. We typically start with an initial wave function as that of a complex Gaussian with a constant velocity: $\exp[-(x^2/2q_x^2 + y^2/2q_y^2 + z^2/2q_z^2) - i\vec{k} \cdot \vec{r}]$. q_x, q_y, q_z , and $\vec{k}$ are adjustable parameters which are checked to ensure that their choices do not affect the final converged ground state [14].

For $c_2 = 0$ or $c = 0$, it is easy to check that SMA is always valid since the energy functional is symmetric with respect to spin component index. The fractional populations for each component is therefore the same as for a homogeneous system, i.e. given by $(\frac{1-n_0+m}{2}, n_0, \frac{1-n_0-m}{2})$ if $B = 0$, and $(m, 1 - m, 0)$ if $B > 0$.

For ^{87}Rb and ^{23}Na condensates, which are believed to be ferromagnetic $c_2 < 0$ ($c < 0$) and anti-ferromagnetic $c_2 > 0$ ($c > 0$) respectively, Figure 5 gives typical density distributions of spacial mode function, $\rho(\vec{r}) = |\phi_j(\vec{r})|^2$. Both sub-figures clearly indicate that SMA is no longer valid. To get an overall idea of the validity of SMA we plot in Figure 6 the overlap integrals of our mode functions with respect to the SMA mode function $\phi_{\text{SMA}}(\vec{r})$ as determined from a scalar GP equation with a nonlinear coefficient $\propto c_0$ (due to the symmetric H_S only) [14]. For a ^{87}Rb condensate, we see the overlap is close to unity when B is small, therefore, SMA remains approximately applicable. But it becomes increasingly bad with the increase of B . We thus conclude that the SMA remains reasonable in a weak magnetic field while it is clearly invalid in a strong B -field. In fact, our numerical results confirm that the stronger the B -field, the




Figure 6. The overlap between the SMA mode function and the mode function for $|-\rangle$ component. Left panel is for a ^{87}Rb condensate. Right panel is for a ^{23}Na condensate. The atomic parameters are the same as in Fig. 5.




Figure 7. The same as in Fig. 6, but now comparing the spin asymmetric energy $E_a = c_2 \langle \vec{F} \rangle^2 / 2 - (\eta_0 + \eta) \langle F_z \rangle + \delta \langle F_z^2 \rangle$ with the spin symmetric one H_s .

worse the SMA gets. For typical system parameters, the dividing line occurs at a B -field of a fraction of a Gauss when the system magnetization $\mathcal{M}$ is not too small or too large. For a condensate with anti-ferromagnetic interactions, it was found earlier that SMA is violated in the limit of both large N and $\mathcal{M}$ even without an external B -field, while the case of $\mathcal{M} = 0$ presents an exception where SMA remains strictly valid for $B = 0$ [14]. Figure 6 shows the overlap integral for a ^{23}Na condensate, indeed we see SMA is invalid except at $\mathcal{M} = 0$ where all atoms are in the $|0\rangle$ component. Remarkably, despite the seemingly large deviations from the SMA (as in Fig. 6), the spin asymmetric energy term remains very small in comparison to the spin symmetric term as evidenced in Fig. 7.

Figure 8 shows the dependence of fractional populations on the fractional magnetization for a ^{87}Rb (left column) and a ^{23}Na condensate (right column) at different B -fields. For ^{87}Rb atoms, these curves resemble the same dependence as for a homogeneous system where SMA is strictly valid. Nevertheless, we find the densities of mode functions can become quite different, i.e. SMA is not valid in general. For




Figure 8. Fractional population for each spin component of a ^{87}Rb (left column) and a ^{23}Na (right column) condensate. The values of B -field from top row to bottom are $B = 0, 0.1, 1.0$ (Gauss). The atomic parameters are the same as in Fig. 5. The solid lines with plus signs denote the $|+\rangle$ component, the lines with triangles are for the $|-\rangle$ component and the lines with open circles for the $|0\rangle$ component. The vertical dashed lines in (e) and (f) indicate the critical value m_c , the boundary between the two distinct regions discussed in the text. In (d), $m_c = 0$.

^{23}Na atoms, the fractional component populations at different B -fields again follow the results as obtained previously for the homogeneous case. When $B = 0$ [as in Fig. 8(d)], the ground state distribution clearly obeys the same earlier (homogeneous) result $(\frac{1+m}{2}, 0, \frac{1-m}{2})$, including the special case when $\mathcal{M} = 0$ where it becomes $(\frac{1-n_0}{2}, n_0, \frac{1-n_0}{2})$ with $n_0 \in [0, 1]$. For $B \neq 0$ [as in Fig. 8(e) and (f)], our numerical solutions reveal again two distinct regions; one for $m < m_c$ where all three components coexist, and another one for $m > m_c$ where only two components ($|+\rangle$ and $|-\rangle$) coexist. We find that m_c increases with the B -field, and is of course limited to $m_c < 1$. We conclude that despite the fact a harmonic trap induces spatially inhomogeneous distribution to condensate density, thus breaks the SMA in general, the overall ground state properties as measured by the fractional component distributions follow closely the results as obtained previously for the homogeneous case. Physically, we believe the above results can be understood as fractional populations relate to integrals of wave functions over all spaces, during which differences between wave functions can be averaged out. When only the $|+\rangle$ and $|-\rangle$ components coexist, in fact, the two constraints on N and $\mathcal{M}$




Figure 9. The B field dependence of the critical fractional magnetization m_c as computed numerically for a ^{23}Na condensate in a harmonic trap. The smooth curve corresponds to the result of $\delta = 2c[1 - \sqrt{1 - m_c^2}]$ (as from the homogeneous case) with an appropriately adjusted coefficient c (or density). The atomic parameters are the same as in Fig. 5.

always give the fractional population $n_{\pm} = (1 \pm m)/2$ if $N_0 = 0$.

5. Conclusion

We have revisited the question of the mean field ground state structures of spin-1 condensate in the presence of a uniform magnetic field. For a homogeneous system, when $c = 0$, there exists in general only two nonzero components $|+\rangle$ and $|0\rangle$, except when $B = 0$ where the ground state solution becomes indefinite; for ferromagnetic interactions when $c < 0$, the ground state in general has three nonzero components; when $c > 0$ as for anti-ferromagnetic interactions, except for $m = 0$, there are two regions: one for $\delta > 2c[1 - \sqrt{1 - m_c^2}]$ where three nonzero components coexist and one for $\delta \leq 2c[1 - \sqrt{1 - m_c^2}]$ where only two components coexist. Inside a harmonic trap, these results remain largely true, although the SMA becomes generally invalid. We find interestingly (see Fig. 9), the B field (or the δ) dependence of the critical value m_c that separates the two and three component condensate regions, remains almost identical as that given by the analytical formulae $\delta = 2c[1 - \sqrt{1 - m_c^2}]$ for a homogeneous system. In a sense, this also points to the validity of the use of a mean field description, as the number of atoms is really large (10^6).

6. Acknowledgement

We acknowledge interesting discussions with Prof. M. S. Chapman and Mr. M. -S. Chang. This work is supported by NSF.

References

- [1] See <http://amo.phy.gasou.edu/bec.html/bibliography.html>
- [2] Anderson M H, Ensher J R, Matthews M R, Wieman C E and Cornell E A 1995 *Science* **269** 198; Davis K B, Mewes M -O, Andrews M R, Druten N J van, Durfee D S, Kurn D M and Ketterle W 1995 *Phys. Rev. Lett.* **75** 3969; Bradley C C, Sackett C A, Tollett J J and Hulet R G 1995 *Phys. Rev. Lett.* **75** 1687; ———1997 *Phys. Rev. Lett.* **79** 1170
- [3] Dalfovo F, Giorgini S, Pitaevskii L P and Stringari S 1999 *Rev. Mod. Phys.* **71** 463
- [4] Stamper-Kurn D M, Andrews M R, Chikkatur A P, Inouye S, Miesner H -J, Stenger J and Ketterle W 1998 *Phys. Rev. Lett.* **80** 2027
- [5] Barrett M D, Sauer J A and Chapman M S 2001 *Phys. Rev. Lett.* **87** 010404
- [6] Stenger J, Inouye S, Stamper-Kurn D M, Miesner H -J, Chikkatur A P and Ketterle W 1998 *Nature* **396** 345
- [7] Ho T -L 1998 *Phys. Rev. Lett.* **81** 742
- [8] Law C K, Pu H, and Bigelow N P 1998 *Phys. Rev. Lett.* **81** 5257
- [9] Pu H, Law C K, and Bigelow N P 2000 *Physica B* **280** 27
- [10] Ohmi T and Machida K 1998 *J. Phys. Soc. Jpn.* **67** 1822
- [11] Pu H 1999 *Phys. Rev. A* **60** 1463
- [12] Vanier J and Audoin C 1988 *The quantum physics of atomic frequency standards* (Philadelphia: A. Hilger)
- [13] Stamper-Kurn D M and Ketterle W 1999 *Spinor condensates and light scattering from Bose-Einstein condensates*, in the Proceedings of Les Houches 1999 Summer School, Session LXXII, (New York: Springer-Verlag).
- [14] Yi S, Müstecaplıoğlu Ö E, Sun C P and You L 2002 *Phys. Rev. A* **66** 011601
- [15] Kempen E G M van, Kokkelmans S J J M F, Heinzen D J and Verhaar B J 2002 *Phys. Rev. Lett.* **88** 093201
- [16] Crubellier A, Dulieu O, Masnou-Seeuws F, Elbs M, Knöckel H and Tiemann E 1999 *Eur. Phys. J. D* **6** 211
- [17] Ho T -L and Yip S K 2000 *Phys. Rev. Lett.* **84** 4031
- [18] Koashi M and Ueda M 2000 *Phys. Rev. Lett.* **84** 1066